

# PROTOCOL FOR DISTINCTNESS, UNIFORMITY AND STABILITY TESTS

Allium porrum L.

**LEEK** 

UPOV Code: ALLIU\_POR

Adopted on 01/04/2009

Entered into force on 12/03/2009

# I - SUBJECT OF THE PROTOCOL

The protocol describes the technical procedures to be followed in order to meet the Council Regulation 2100/94 on Community Plant Variety Rights. The technical procedures have been agreed by the Administrative Council and are based on general UPOV Document TG/1/3 and UPOV Guideline TG/085/7 dated 09/04/2008 for the conduct of tests for Distinctness, Uniformity and Stability. This protocol applies to varieties of *Allium porrum* L.

### II - SUBMISSION OF SEED AND OTHER PLANT MATERIAL

- 1. The Community Plant Variety Office (CPVO) is responsible for informing the applicant of
  - the closing date for the receipt of plant material;
  - the minimum amount and quality of plant material required;
  - the examination office to which material is to be sent.

A sub-sample of the material submitted for test will be held in the variety collection as the definitive sample of the candidate variety.

The applicant is responsible for ensuring compliance with any customs and plant health requirements.

#### 2. Final dates for receipt of documentation and material by the Examination Office

The final dates for receipt of requests, technical questionnaires and the final date or submission period for plant material will be decided by the CPVO and each Examination Office chosen.

The Examination Office is responsible for immediately acknowledging the receipt of requests for testing, and technical questionnaires. Immediately after the closing date for the receipt of plant material the Examination Office should inform the CPVO whether acceptable plant material has been received or not. However if unsatisfactory plant material is submitted the CPVO should be informed as soon as possible.

#### 3. Plant material requirements

The current quality and quantity requirements as well as the final dates for submission of the plant material are available on the CPVO website (<a href="www.cpvo.europa.eu">www.cpvo.europa.eu</a>) and are published in the CPVO gazette 'S2'.

Labelling of sample: ..... - Species

- File number of the application allocated by the CPVO
- Breeder's reference
- Examination reference (if known)
- Name of applicant
- The phrase "On request of the CPVO"

#### III - CONDUCT OF TESTS

#### 1. Variety collection

A variety collection will be maintained for the purpose of establishing distinctness of the candidate varieties in test. A variety collection may contain both living material and descriptive information. A variety will be included in a variety collection only if plant material is available to make a technical examination.

Pursuant to Article 7 of Council Regulation No. 2100/94, the basis for a collection should be the following:

- varieties listed or protected at the EU level or at least in one of the EEA Member States;
- varieties protected in other UPOV Member States;
- any other variety in common knowledge.

The composition of the variety collection in each Examination Office depends on the environmental conditions in which the Examination Office is located.

Variety collections will be held under conditions which ensure the long term maintenance of each accession. It is the responsibility of Examination Offices to replace reference material which has deteriorated or become depleted. Replacement material can only be introduced if appropriate tests confirm conformity with the existing reference material. If any difficulties arise for the replacement of reference material Examination Offices must inform the CPVO. If authentic plant material of a variety cannot be supplied to an Examination Office the variety will be removed from the variety collection.

#### 2. Material to be examined

Candidate varieties will be directly compared with other candidates for Community plant variety rights tested at the same Examination Office, and with appropriate varieties in the variety collection. When necessary an Examination Office may also include other candidates and varieties. Examination Offices should therefore make efforts to coordinate the work with other Offices involved in DUS testing of leek. There should be at least an exchange of technical questionnaires for each candidate variety, and during the test period, Examination Offices should notify each other and the CPVO of candidate varieties which are likely to present problems in establishing distinctness. In order to solve particular problems Examination Offices may exchange plant material.

### 3. Characteristics to be used

The characteristics to be used in DUS tests and preparation of descriptions shall be those referred to in the Annex 1. All the characteristics shall be used, providing that observation of a characteristic is not rendered impossible by the expression of any other characteristic, or the expression of a characteristic is prevented by the environmental conditions under which the test is conducted. In the latter case, the CPVO should be informed. In addition the existence of some other regulation e.g. plant health, may make the observation of the characteristic impossible.

The Administrative Council empowers the President, in accordance with Article 23 of Commission Regulation N° 1239/95, to insert additional characteristics and their expressions in respect of a variety.

#### 4. Grouping of varieties

The varieties and candidates to be compared will be divided into groups to facilitate the assessment of distinctness. Characteristics which are suitable for grouping purposes are those which are known from experience not to vary, or to vary only slightly, within a variety and which in their various states of expression are fairly evenly distributed throughout the collection. In the case of continuous grouping characteristics overlapping states of expression between adjacent groups is required to reduce the risks of incorrect allocation of candidates to groups. The characteristics used for grouping could be the following:

a) Leaf blade: width (characteristic 5)

b) Leaf blade: colour (characteristic 6)

c) Plant: length (characteristic 10)

d) Shaft: length (characteristic 11)

e) Only for vegetatively propagated varieties: Flower: male sterility (characteristic 17)

#### 5. <u>Trial designs and growing conditions</u>

The minimum duration of tests will normally be two independent growing cycles. For vegetatively propagated varieties, the duration of the testing may be reduced to one growing cycle if the results on distinctness and uniformity are conclusive. Tests will be carried out under conditions ensuring normal growth. The size of the plots will be such that plants or parts of plants may be removed for measuring and counting without prejudice to the observations which must be made up to the end of the growing period.

#### The test design is as follows:

As a minimum, each test should include a total of 200 plants for seed-propagated varieties, or 60 plants for vegetatively propagated varieties, divided between two or more replicates.

All observations determined by measurement or counting should be made on 60 plants or parts of 60 plants for seed-propagated varieties, and 20 plants or parts of 20 plants for vegetatively propagated varieties.

#### Special tests

In accordance with Article 83(3) of Council Regulation No. 2100/94 an applicant may claim either in the Technical Questionnaire or during the test that a candidate has a characteristic which would be helpful in establishing distinctness. If such a claim is made and is supported by reliable technical data, a special test may be undertaken providing that a technically acceptable test procedure can be devised.

Special tests will be undertaken, with the agreement of the President of CPVO, where distinctness is unlikely to be shown using the characters listed in the protocol.

#### 7. Standards for decisions

### a) Distinctness

A candidate variety will be considered to be distinct if it meets the requirements of Article 7 of Council Regulation No. 2100/94.

#### b) Uniformity

For the assessment of uniformity of:

- (i) cross pollinated varieties: relative uniformity standards should be applied.
- (ii) single cross hybrids: a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 200 plants, 7 off-types are allowed.

(iii) – vegetatively propagated varieties: a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 60 plants, 2 off-types are allowed.

### c) Stability

A candidate will be considered to be sufficiently stable when there is no evidence to indicate that it lacks uniformity.

#### IV - REPORTING OF RESULTS

After each recording season the results will be summarised and reported to the CPVO in the form of a UPOV model interim report in which any problems will be indicated under the headings distinctness, uniformity and stability. Candidates may meet the DUS standards after two growing periods but in some cases three growing periods may be required. When tests are completed the results will be sent by the Examination Office to the CPVO in the form of a UPOV model final report.

If it is considered that the candidate complies with the DUS standards, the final report will be accompanied by a variety description in the format recommended by UPOV. If not the reasons for failure and a summary of the test results will be included with the final report.

The CPVO must receive interim reports and final reports by the date agreed between the CPVO and the examination office.

Interim reports and final examination reports shall be signed by the responsible member of the staff of the Examination Office and shall expressly acknowledge the exclusive rights of disposal of CPVO.

#### V - LIAISON WITH THE APPLICANT

If problems arise during the course of the test the CPVO should be informed immediately so that the information can be passed on to the applicant. Subject to prior agreement, the applicant may be directly informed at the same time as the CPVO particularly if a visit to the trial is advisable.

The interim report as well as the final report shall be sent by the Examination Office to the CPVO.

#### VI - ENTRY INTO FORCE

The present protocol enters into force on **12 March 2009**. Any ongoing DUS examination of candidate varieties started before the aforesaid date will not be affected by the approval of the new TP. Technical examinations of candidate varieties are carried out according to the TP in force when the DUS test starts. The starting date of a DUS examination is considered to be the due date for the submission of plant material for the first growing period.

In cases where the CPVO requests to take-over a DUS report for which the technical examination has either been finalized or which is in the process of being carried out at the moment of the request, such report can only be accepted if the technical examination has been carried out according to the CPVO TP which was in force at the moment when the technical examination started.

\*\*\*\*\*

# **ANNEXES TO FOLLOW**

| ANNEX I   | <u>PAGE</u>   |
|---|---|
| Table of characteristics  | 7   |
| Explanations and methods  | 11  |
| <u>Legend</u> :   |   |
| Note: For the CPVO numbered characteristics, all characteristics in the table are compuls the case of disease resistance characteristics, only those resistances marked with an asterisk (*) is compulsory. The asterisks in the UPOV numbered characteristics are there for information purpocharacteristics which should always be observed when a UPOV guideline is utilised. In general for the assessment of resistance characteristics, the facilities of other Examination Offices of might be used, subject to previous arrangements.  Some characteristics may be discarded: if there are already phytosanitary restrictions.  | in the CPVO column are poses and denote those   |
| (+) See explanations on the Table of characteristics (a) – (c) See explanations on the table of characteristics Grouping characteristic   |   |
| Types of expression of characteristics:   |   |
| QL – Qualitative characteristic<br>QN – Quantitative characteristic<br>PQ – Pseudo-qualitative characteristic   |   |
| Type of observation of characteristics:   |   |
| MG – Single measurement of a group of plants or parts of plants<br>MS – Measurement of a number of individual plants or parts of plants<br>VG – Visual assessment by a single observation of a group of plants or parts of plants<br>VS – Visual assessment by observation of individual plants or parts of plants  |   |
| When a method of observation is attributed to a certain characteristic, the first differentiation is maction taken is a <u>visual observation (V)</u> or a <u>measurement (M).</u>  | nade depending if the   |
| The second differentiation deals with the number of observations the expert attributes to each variattribution of either G or S.  If a single observation of a group consisting of an undefined number of individual plants is approprexpression of a variety, we talk about a visual observation or a measurement made on a group of attribute the letter G (either VG or MG). If the expert makes more than one observation on that g decisive part is that we have at the end only one data entry per variety which means that we have measurement of plant length on a plot – MG, visual observation of green colour of leaves on a plot it is necessary to observe a number of individual plants to assess the expression of a variety, we letter S (thus either VS or MS). Single plant data entries are kept per variety for further calculation (e.g. measurement of length of ears – MS, visual observation of growth habit of single plants in g number of individual plants to be observed in such cases is stated in section III.5. | priate to assess the plants, thus we proup of plants, the e to deal with G (e.g. ot – VG).  e should attribute the ns like the variety mean |
| Literature  | 16  |

# **ANNEX II**

**Technical Questionnaire** 

ANNEX I

TABLE OF CHARACTERISTICS TO BE USED IN DUS-TEST AND PREPARATION OF DESCRIPTIONS

| CPVO<br>No. | UPOV<br>No. | Stage,<br>Method | Characteristics               | Examples  | Note |
|-------------|-------------|------------------|-------------------------------|---|------|
| 1.          | 1.          | VG/MS            | Plant: height                 |   |      |
| (+)         | (+)         | (a)              | very short                    | De Carentan 2                                       | 1    |
| QN          | QN          |                  | short D'Hiver de Saint Victor |   | 3    |
|             |             |                  | medium Bleu de Solaise        |   | 5    |
|             |             |                  | tall                          | Kingston, Long géant du Verdet                      | 7    |
|             |             |                  | very tall                     | Bulgaarse Reuzen                                    | 9    |
| 2.          | 2.          | VG               | Foliage: attitude             |   |      |
| (+)         | (+)         | (a)              | erect                         | Rese  | 1    |
| QN          | QN          |                  | semi-erect                    | Linx, Upton   | 3    |
|             |             |                  | horizontal                    | De Carentan 2, D'Elbeuf,<br>Jaune gros du Poitou    | 5    |
| 3.          | 3.          | VG               | Leaf blade: bending           |   |      |
| (+)         | (+)         | (b)              | weak                          | Bell, Lampton                                       | 3    |
| QN          | QN          |                  | medium                        | Flextan, Linx                                       | 5    |
|             |             |                  | strong                        | Blauwgroene Winter, Bulgaarse<br>Reuzen             | 7    |
| 4.          | 4.          | VG/MS            | Leaf blade: length            |   |      |
| (+)         | (+)         | (b)              | short                         | Artemis, Conora, De Carentan 2,<br>D'Elbeuf, Kalmar | 3    |
| QN          | QN          |                  | medium                        | Flextan, Porridor, Rese                             | 5    |
|             |             |                  | long                          | Kingston, Kong Richard, Maxim                       | 7    |
| 5.          | 5.<br>(*)   | VG/MS            | Leaf blade: width             |   |      |
| (+)         | (+)         | (b)              | narrow                        | Lampton, Rustic                                     | 3    |
| QN          | QN          |                  | medium                        | De Liège  | 5    |
| G           |             |                  | broad                         | Jaune gros du Poitou, Rese, Striker                 | 7    |

| CPVO<br>No. | UPOV<br>No. | Stage,<br>Method | Characteristics   | Examples                                 | Note |
|-------------|-------------|------------------|---|--|------|
| 6.          | 6.          | VG               | Leaf blade: colour                                      |  |      |
|             | (*)         | (b)              | yellow green  | Jaune gros du Poitou                     | 1    |
| PQ          | PQ          |                  | green   | Premier                                  | 2    |
|             |             |                  | grey green  | Zwitserse Reuzen                         | 3    |
| G           |             |                  | blue green Blauwgroene Winter, Libertas, Olaf, Porridor |  | 4    |
| 7.          | 7.          | VG               | Leaf blade: intensity of colour                         |  |      |
| QN          | QN          | (b)              | light   |  | 3    |
|             |             |                  | medium  |  | 5    |
|             |             |                  | dark  |  | 7    |
| 8.          | 8.          | VG               | Leaf blade: anthocyanin coloration                      |  |      |
| (+)         | (+)         | (b)              | absent or very weak                                     | Jaune gros du Poitou                     | 1    |
| QN          | QN          |                  | weak  | Azur                                     | 3    |
|             |             |                  | medium  |  | 5    |
|             |             |                  | strong  | Nepal                                    | 7    |
|             |             |                  | very strong   | D'Hiver de Saint Victor                  | 9    |
| 9.          | 9.          | VG               | Leaf blade: waxiness                                    |  |      |
| QN          | QN          | (b)              | absent or very weak                                     | Jaune gros du Poitou, Kingston           | 1    |
|             |             |                  | weak  | Carlton, Gros long d'été 2, Rese         | 3    |
|             |             |                  | medium  | D'Elbeuf, Linx                           | 5    |
|             |             |                  | strong  | Bleu de Solaise, Flextan                 | 7    |
|             |             |                  | very strong   |  | 9    |
| 10.         | 10.         | VG/MS            | Plant: length   |  |      |
|             | (*)         | (c)              | very short  |  | 1    |
| (+)         | (+)         |                  | short   | De Carentan 2                            | 3    |
| QN          | QN          |                  | medium  | Bleu de Solaise,<br>Jaune gros du Poitou | 5    |
|             |             |                  | long  | Kingston, Rese, Titan                    | 7    |
| G           |             |                  | very long   | Bulgaarse Reuzen                         | 9    |

| CPVO<br>No. | UPOV<br>No. | Stage,<br>Method | Characteristics               | Examples   | Note |
|-------------|-------------|------------------|-------------------------------|--|------|
| 11.         | 11.         | VG/MS            | Shaft: length                 |  |      |
|             | (*)         | (c)              | very short                    |  | 1    |
| (+)         | (+)         |                  | short                         | Artemis, Bleu de Solaise,<br>D'Hiver de Saint Victor | 3    |
| QN          | QN          |                  | medium                        | Gros long d'été 2                                    | 5    |
|             |             |                  | long                          | Lampton, Maxim                                       | 7    |
| G           |             |                  | very long                     | Kingston Kong, Richard                               | 9    |
| 12.         | 12.         | VG/MS            | Shaft: diameter               |  |      |
|             | (*)         | (c)              | very small                    |  | 1    |
| (+)         | (+)         |                  | small                         | Lampton, Titan                                       | 3    |
| QN          | QN          |                  | medium                        | Géant précoce  | 5    |
|             |             |                  | large                         | Premier, Zwitserse Reuzen                            | 7    |
|             |             |                  | very large                    | Jaune gros du Poitou                                 | 9    |
| 13.         | 13.         | VG               | Shaft: ratio length/diameter  |  |      |
| (+)         | (+)         | (c)              | small                         | D'Hiver de Saint Victor                              | 3    |
| QN          | QN          |                  | medium                        | Easton, Gros long d'été 2                            | 5    |
|             |             |                  | large                         | Bulgaarse Reuzen                                     | 7    |
| 14.         | 14.         | VG               | Shaft: bulb formation         |  |      |
|             | (*)         | (c)              | absent or very weak           | Jolant, Roxton, Striker                              | 1    |
| QN          | QN          |                  | weak                          | Hafnia, Lampton, Linx, Titan                         | 3    |
|             |             |                  | medium                        | Bleu de Solaise, Premier                             | 5    |
|             |             |                  | strong                        | Artemis, Jaune gros du Poitou                        | 7    |
|             |             |                  | very strong                   | Carentan 2   | 9    |
| 15.         | 15.         | VG               | Shaft: narrowing towards base |  |      |
| (+)         | (+)         | (c)              | absent                        | Herfstreuzen 2                                       | 1    |
| QL          | QL          |                  | present                       | D'Elbeuf   | 9    |

| CPVO<br>No. | UPOV<br>No. | Stage,<br>Method | Characteristics  | Examples | Note |
|-------------|-------------|------------------|--|----------|------|
| 16.         | 16.         | VG               | Only for vegetatively propagated varieties: Spathe: length         |          |      |
| (+)         | (+)         |                  | short  |          | 3    |
| QN          | QN          |                  | medium   |          | 5    |
|             |             |                  | long   |          | 7    |
| 17.         | 17.<br>(*)  | VG               | Only for vegetatively propagated varieties: Flower: male sterility |          |      |
| (+)         | (+)         |                  | absent   |          | 1    |
| QL          | QL          |                  | present  |          | 9    |
| G           |             |                  |  |          |      |


# **EXPLANATIONS AND METHODS**

# **Explanations for covering several characteristics**


Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Observations should be made when the outer (oldest) leaf blades start to senesce.
- (b) Observations on the leaf blade should be made on the fully developed leaf.
- (c) Observations should be made at harvest maturity.


# Ad 1: Plant: height



Ad 2: Foliage: attitude




Ad. 3: Leaf blade: bending



Ad 4: Leaf blade: length Ad 5: Leaf blade: width

Leaf blade length and width should be observed on the longest leaf blade.



# Ad 8: Leaf blade: anthocyanin coloration

To be observed after a period of night frost.


# Ad 10: Plant: length

Plant length should be measured on the harvested plant with the longest leaves brought in line with the shaft.




Ad 11: Shaft: length

Ad 11: Shaft: diamter
Ad 11: Shaft: ratio length/diameter



The diameter of the shaft should be observed at the middle of the length.

# Ad 13: Shaft: ration length/diameter




Ad 15: Shaft: narrowing towards base



# Ad 16: Spathe: length

To be observed on the green, fully developed spathe, before it starts to open and desiccate.



# Ad 17: Flower: male sterility

The observation of male sterility should be done at full flowering. Male sterility can be observed by rubbing a piece of black paper gently over the flower head: if the flowers are sterile, no pollen will be seen on the paper. In addition, in the case of male sterile varieties, the anthers are empty and desiccate very quickly. Further explanation required?

# **LITERATURE**

Bonnet, B., 1976: Le poireau (*Allium porrum* L.): Aspects botaniques et agronomiques. Revue bibliographique, Saussurea 7, pp. 175-194.

Currah, L., 1986: Leek breeding: A review. Review Article, Journal of Horticultural Science 61, (4) 407-415.

Jones, H.A., Mann, L.K., 1963: Onions and Their Allies: Botany, Cultivation and Utilisation, London, GB, Leonard Hill.

Rabinowitch, H. D., Brewster, J. L., 1990: Onions and Allied Crops. Vol. 1-3, CRC Press, Boca Raton.

Schweisguth, B., 1970: Études préliminaires à l'amélioration du poireau *A. porrum* L. Proposition d'une méthode d'amélioration, Annales de l'Amélioration des Plantes 20, pp. 215-231.

Schweisguth B., 1973: Étude de l'hérédité de trois caractères quantitatifs chez le poireau (*Allium porrum* L.). Annales de l'Amélioration des Plantes 23, pp. 45-57.

# **ANNEX II**



|    | TECHNICAL QUESTIONNAIRE  |
|----|--|
|    | to be completed in connection with an application for Community Plant Variety Rights Please answer all questions. A question without any answer will lead to a non-attribution of an application date. In cases where a field / question is not applicable, please state so. |
| 1. | Botanical taxon: Name of the genus, species or sub-species to which the variety belongs and common name  |
|    | Allium porrum L.   |
|    | LEEK   |
|    |  |
| 2. | <b>Applicant(s):</b> Name(s) and address(es), phone and fax number(s), Email address, and where appropriate name and address of the procedural representative  |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |
| 3. | Variety denomination   |
|    | a) Where appropriate proposal for a variety denomination:  |
|    |  |
|    | b) Provisional designation (breeder's reference):  |
|    |  |
|    |  |

| Information on origin, maintenance and reproduction of the variety |                               |   |   |  |  |  |  |  |
|--|-------------------------------|---|---|--|--|--|--|--|
|  |                               |   |   |  |  |  |  |  |
| (a)  | (i) hybrid                    |   |   |  |  |  |  |  |
|  |                               |   |   |  |  |  |  |  |
|  |                               |   |   |  |  |  |  |  |
|  | (III) par                     | rent line   |   |  |  |  |  |  |
| (b)  | (i) seed                      | d propagated[ ]   |   |  |  |  |  |  |
|  | (ii) veg                      | etatively propagated[ ]   |   |  |  |  |  |  |
| (c)  | Other i                       | information on genetic origin and breeding method[ ]  |   |  |  |  |  |  |
| Method   | of propa                      | ngation   |   |  |  |  |  |  |
| (  | a) See                        | ed propagated varieties   |   |  |  |  |  |  |
|  | (i)                           | Cross-pollination   |   |  |  |  |  |  |
|  | (ii)                          | Hybrid  |   |  |  |  |  |  |
|  |                               | seed-propagated parents[ ]  |   |  |  |  |  |  |
|  |                               |   |   |  |  |  |  |  |
|  |                               | one vegetatively propagated and one seed-propagated parent  |   |  |  |  |  |  |
|  |                               | one vegetatively propagated and one seed-propagated parent  |   |  |  |  |  |  |
|  | (iii)                         |   |   |  |  |  |  |  |
| (  |                               | two vegetatively propagated parents   |   |  |  |  |  |  |
| (  |                               | two vegetatively propagated parents   |   |  |  |  |  |  |
| (  | b) Ve                         | two vegetatively propagated parents   |   |  |  |  |  |  |
| (  | b) Ve                         | two vegetatively propagated parents   |   |  |  |  |  |  |
| (  | b) Ver                        | two vegetatively propagated parents   |   |  |  |  |  |  |
|  | b) Ve<br>(i)<br>(ii)<br>(iii) | two vegetatively propagated parents   | ered  |  |  |  |  |  |
| Geogra   | b) Ve<br>(i)<br>(ii)<br>(iii) | two vegetatively propagated parents   | ered  |  |  |  |  |  |
| Geogra   | b) Ve<br>(i)<br>(ii)<br>(iii) | two vegetatively propagated parents   | ered  |  |  |  |  |  |
|  | Bree Pleas (a) (b) (c)        | Breeding, m Please indicat  (a)  (i) hybriding  (ii) oper  (iii) par  (b)  (i) seed  (ii) veg  (c) Other   Method of proparion  (a) Seed  (i) | Breeding, maintenance and reproduction of the variety Please indicate breeding scheme, parents, other relevant information  (a)  (i) hybrid |  |  |  |  |  |

| 4.4        | Shall the information on data relating to components of hybrid varieties including data related to their cultivation be treated as confidential? |   |                         |  |  |  |  |  |
|------------|--|---|-------------------------|--|--|--|--|--|
|            | [ ] YES [ ] NO   |   |                         |  |  |  |  |  |
|            | If yes, please give this information on the  | If yes, please give this information on the attached form for confidential information.                   |                         |  |  |  |  |  |
|            | If no, please give information on data relating to components of hybrid varieties including data related to their cultivation:                   |   |                         |  |  |  |  |  |
|            | Breeding scheme (indicate female compo   | nent first)   |                         |  |  |  |  |  |
|            |  |   |                         |  |  |  |  |  |
|            |  |   |                         |  |  |  |  |  |
|            |  |   |                         |  |  |  |  |  |
| 5.         |  | <b>ndicated</b> (the number in brackets refers to the Protocol; please mark the state of expression which | best                    |  |  |  |  |  |
|            | Characteristics Example varieties Note   |   |                         |  |  |  |  |  |
|            | Characteristics  | Example varieties   | Note                    |  |  |  |  |  |
| 5.1<br>(5) | Characteristics  Leaf blade: width   | Example varieties   | Note                    |  |  |  |  |  |
|            |  | Example varieties  Lampton, Rustic  | Note 3 [ ]              |  |  |  |  |  |
|            | Leaf blade: width  | ·   |                         |  |  |  |  |  |
|            | Leaf blade: width  | Lampton, Rustic   | 3[]                     |  |  |  |  |  |
|            | Leaf blade: width  narrow  medium  | Lampton, Rustic  De Liège   | 3 [ ]<br>5 [ ]          |  |  |  |  |  |
| 5.2        | Leaf blade: width  narrow  medium  broad   | Lampton, Rustic  De Liège   | 3 [ ]<br>5 [ ]          |  |  |  |  |  |
| 5.2        | Leaf blade: width  narrow  medium  broad  Leaf blade: colour   | Lampton, Rustic  De Liège  Jaune gros du Poitou, Rese, Striker  | 3 [ ]<br>5 [ ]<br>7 [ ] |  |  |  |  |  |
| 5.2        | Leaf blade: width  narrow  medium  broad  Leaf blade: colour  yellow green   | Lampton, Rustic  De Liège  Jaune gros du Poitou, Rese, Striker  Jaune gros du Poitou                      | 3 [ ]<br>5 [ ]<br>7 [ ] |  |  |  |  |  |

|             | Characteristics             | Example varieties                                    | Note  |
|-------------|-----------------------------|--|-------|
| 5.3<br>(10) | Plant: length               |  |       |
|             | very short                  |  | 1[]   |
|             | short                       | De Carentan 2  | 3 [ ] |
|             | medium                      | Bleu de Solaise, Jaune gros du Poitou                | 5 [ ] |
|             | long                        | Kingston, Rese, Titan                                | 7 [ ] |
|             | very long                   | Bulgaarse Reuzen                                     | 9[]   |
| 5.4         | Shaft: length               |  |       |
| (11)        | very short                  |  | 1[]   |
|             | short                       | Artemis, Bleu de Solaise,<br>D'Hiver de Saint Victor | 3 [ ] |
|             | medium                      | Gros long d'été 2                                    | 5 [ ] |
|             | long                        | Lampton, Maxim                                       | 7 [ ] |
|             | very long                   | Kingston Kong, Richard                               | 9[]   |
| 5.5<br>(14) | Shaft: bulb formation       |  |       |
|             | absent or very weak         | Jolant, Roxton, Striker                              | 1[]   |
|             | weak                        | Hafnia, Lampton, Linx, Titan                         | 3 [ ] |
|             | medium                      | Bleu de Solaise, Premier                             | 5 [ ] |
|             | strong                      | Artemis, Jaune gros du Poitou                        | 7 [ ] |
|             | very strong                 | Carentan 2   | 9[]   |
| 5.6<br>(17) | Only for vegetatively propa | agated varieties: Flower: male sterility             |       |
|             | absent                      |  | 1[]   |
|             | present                     |  | 9[]   |

| 6.  | Similar varieties and differences from these varieties: |  |  |  |  |  |
|-----|---|--|--|--|--|--|
|     | Denomination of similar variety                         | Characteristic in which the similar variety is different <sup>1)</sup> | State of expression of similar variety | State of expression of candidate variety |  |  |
|     |   |  |  |  |  |  |
|     |   |  |  |  |  |  |
| 1)  |   |  |  |  |  |  |
| 1)  |   | states of expressions of both varie                                    |  | e of the difference                      |  |  |
| 7.  |   | tion which may help to distingu  | ish the variety                        |  |  |  |
| 7.1 | Resistance to pest                                      | s and diseases   |  |  |  |  |
|     |   |  |  |  |  |  |
|     |   |  |  |  |  |  |
| 7.2 | Special conditions                                      | for the examination of the va  | riety                                  |  |  |  |
|     | [ ] YES, please sp                                      | ecify  |  |  |  |  |
|     | [ ] TES, piedse sp                                      | cony   |  |  |  |  |
|     |   |  |  |  |  |  |
|     | [ ] NO  |  |  |  |  |  |
| 7.3 | Other information                                       |  |  |  |  |  |
|     |   |  |  |  |  |  |
|     | [ ] YES, please sp                                      | ecify  |  |  |  |  |
|     |   |  |  |  |  |  |
|     | [ ] NO  |  |  |  |  |  |
|     |   |  |  |  |  |  |
| 7.4 | Growing season  |  |  |  |  |  |
|     | a)  | spring [ ]   |  |  |  |  |
|     | b)  | summer [ ]   |  |  |  |  |
|     | c)  | autumn [ ]   |  |  |  |  |
|     | d)  | winter [ ]   |  |  |  |  |

| 8. | GMO-information required   |                               |                         |        |  |  |
|----|--|-------------------------------|-------------------------|--------|--|--|
|    | The variety represents a Genetically Modified Organism within the meaning of Article 2(2) of Council Directive EC/2001/18 of 12/03/2001.   |                               |                         |        |  |  |
|    | [ ] YES [ ] NO   |                               |                         |        |  |  |
|    | If yes, please add a copy of the written examination of the variety under Article environment according to the norms of the second seco | s 55 and 56 of the Basic Regu | ulation does not pose   |        |  |  |
| 9. | Information on plant material to be  | examined                      |                         |        |  |  |
|    | <b>9.1</b> The expression of a characteristic o such as pests and disease, chemical treculture, different rootstocks, scions take  | atment (e.g. growth retardan  | ts or pesticides), effe |        |  |  |
|    | <b>9.2</b> The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:   |                               |                         |        |  |  |
|    | (a) Microorganisms (e.g. virus, bacteria,  | phytoplasma)                  | [ ] Yes                 | [ ] No |  |  |
|    | (b) Chemical treatment (e.g. growth reta   | ardant or pesticide)          | [ ] Yes                 | [ ] No |  |  |
|    | (c) Tissue culture   |                               | [ ] Yes                 | [ ] No |  |  |
|    | (d) Other factors  |                               | [ ] Yes                 | [ ] No |  |  |
|    | Please provide details of where you have indicated "Yes":  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    | I/we hereby declare that to the best of my/our knowledge the information given in this form is complete and correct.   |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    |  |                               |                         |        |  |  |
|    | Date   | Signature                     | Namo                    | е      |  |  |

[End of document]